

## Lecture 2

Lecturer: Ola Svensson

Scribes: Niklas Gehlen, Tobias Räderscheidt

## 1 Last Lecture

In the previous lecture, we introduced the *probabilistic method*, which allowed us to prove the existence of a combinatorial object with specified properties. In general, the argument goes as follows: select an object at random from the set, and calculate the probability that it satisfies the required property; if this probability is strictly positive then such an object must exist. However this approach is not powerful enough when the events we are considering are not independent.

## 2 Lovász Local Lemma

**Definition 1** An event  $A$  is mutually independent from a set  $\{B_i\}$  of events if for every subset  $\beta$  of events or their complements contained in  $\{B_i\}$

$$\Pr[A \mid \beta] = \Pr[A]$$

In a general setting, we have a set of  $n$  bad events  $\{A_i\}$  that we are trying to avoid, such that  $\Pr[A_i] \leq p < 1$ , for  $i \in \{1, 2, \dots, n\}$ . If we assume that these events are independent, then their complements are independent as well and we can show that

$$\Pr\left[\bigwedge_i \bar{A}_i\right] \geq (1-p)^n > 0$$

However, if we remove the independence assumption, the union bound yields

$$\Pr\left[\bigwedge_i \bar{A}_i\right] \geq 1 - \sum_i \Pr[\bar{A}_i]$$

The *Lovász Local Lemma* improves upon the union bound in the case where the events are not *mutually independent*, but their dependencies are restricted. It was proved by Erdős and Lovász in 1975 [1].

**Theorem 2** (*Lovász Local Lemma*) Let  $A_1, A_2, \dots, A_n$  be a set of "bad" events with  $\Pr[A_i] \leq p < 1$ , and each  $A_i$  is dependent on at most  $d$  other  $A_j$ . If  $p \cdot (d+1) \cdot e \leq 1$ , then

$$\Pr\left[\bigwedge_{i=1}^n \bar{A}_i\right] > 0$$

Before proving the theorem, we give an example showing that the bound is almost tight. Consider events  $A_1, \dots, A_{d+1}$ , each happening with probability  $\frac{1}{d+1}$ . We assume that in each outcome exactly one bad event happens. Consider figure 1. In that case we have  $p \cdot (d+1) = 1$  and  $\Pr\left[\bigwedge_{i=1}^n \bar{A}_i\right] = 0$ .

$A_1$	$A_2$	$A_3$	$A_4$	$A_5$	$A_6$	$A_7$	$A_8$	$A_9$	$A_{10}$
$\frac{1}{10}$	$\frac{1}{10}$	$\frac{1}{10}$	$\frac{1}{10}$	$\frac{1}{10}$	$\frac{1}{10}$	$\frac{1}{10}$	$\frac{1}{10}$	$\frac{1}{10}$	$\frac{1}{10}$

**Figure 1:** Example with  $p = \Pr[A_i] = \frac{1}{10}$  and  $d = 9$

This shows the tightness of the bound. We continue with the proof of the Lovász Local Lemma. To do so, we will use the following lemma:

**Lemma 3** For any  $S \subset \{1, \dots, n\}$  and  $i \in \{1, \dots, n\}$  the following holds:

$$\Pr \left[ A_i \mid \bigwedge_{j \in S} \bar{A}_j \right] \leq \frac{1}{d+1}$$

**Proof** We show the lemma by induction on the size  $m$  of  $S$ .

**Base Case**

For  $m = 0$  we obtain

$$\Pr[A_i] \leq p \leq \frac{1}{(d+1) \cdot e} < \frac{1}{(d+1)}$$

where the second inequation follows by the condition of Lovász Local Lemma.

**Inductive Case**

Assume that the claim is true for all  $S$  with  $|S| < m$ . We prove the claim for an  $S$  with  $|S| = m$ . At first we partition  $S$  into  $S = S_1 \cup S_2$  such that  $S_1$  contains those events of  $S$  that  $A_i$  depends on. Recall that

$$\Pr[A|B] = \frac{\Pr[A \wedge B]}{\Pr[B]}$$

In the following we will apply this equation several times. Using this equation we get

$$\Pr \left[ A_i \mid \bigwedge_{j \in S} \bar{A}_j \right] = \frac{\Pr \left[ A_i \wedge \bigwedge_{j \in S_1} \bar{A}_j \mid \bigwedge_{j \in S_2} \bar{A}_j \right]}{\Pr \left[ \bigwedge_{j \in S_1} \bar{A}_j \mid \bigwedge_{j \in S_2} \bar{A}_j \right]}$$

To bound this term further, we want to give an upper bound on the numerator and a lower bound on the denominator. The upper bound on the numerator is easily obtained as:

$$\Pr \left[ A_i \wedge \bigwedge_{j \in S_1} \bar{A}_j \mid \bigwedge_{j \in S_2} \bar{A}_j \right] \leq \Pr \left[ A_i \mid \bigwedge_{j \in S_2} \bar{A}_j \right] = \Pr[A_i]$$

We continue with the lower bound on the denominator. W.l.o.g. let  $S_1 = \{1, \dots, r\}$ . Therefore, we can rewrite the denominator as:

$$\begin{aligned}
\Pr \left[ \bigwedge_{j=1}^r \bar{A}_j \mid \bigwedge_{j \in S_2} \bar{A}_j \right] &= \Pr \left[ \bar{A}_1 \mid \bar{A}_2 \wedge \bar{A}_3 \wedge \dots \wedge \bar{A}_r \wedge \bigwedge_{j \in S_2} \bar{A}_j \right] \\
&\quad \cdot \Pr \left[ \bar{A}_2 \mid \bar{A}_3 \wedge \bar{A}_4 \wedge \dots \wedge \bar{A}_r \wedge \bigwedge_{j \in S_2} \bar{A}_j \right] \\
&\quad \vdots \\
&\quad \cdot \Pr \left[ \bar{A}_r \mid \bigwedge_{j \in S_2} \bar{A}_j \right] \\
&= \left( 1 - \Pr \left[ A_1 \mid \bar{A}_2 \wedge \bar{A}_3 \wedge \dots \wedge \bar{A}_r \wedge \bigwedge_{j \in S_2} \bar{A}_j \right] \right) \\
&\quad \cdot \left( 1 - \Pr \left[ A_2 \mid \bar{A}_3 \wedge \bar{A}_4 \wedge \dots \wedge \bar{A}_r \wedge \bigwedge_{j \in S_2} \bar{A}_j \right] \right) \\
&\quad \vdots \\
&\quad \cdot \left( 1 - \Pr \left[ A_r \mid \bigwedge_{j \in S_2} \bar{A}_j \right] \right) \\
&\stackrel{I.B.}{\geq} \left( 1 - \frac{1}{d+1} \right) \cdot \left( 1 - \frac{1}{d+1} \right) \cdot \dots \cdot \left( 1 - \frac{1}{d+1} \right) \\
&= \left( 1 - \frac{1}{d+1} \right)^r \geq \left( 1 - \frac{1}{d+1} \right)^d > \frac{1}{e}
\end{aligned}$$

We combine both bounds and obtain:

$$\begin{aligned}
\Pr \left[ A_i \mid \bigwedge_{j \in S} \bar{A}_j \right] &= \frac{\Pr \left[ A_i \wedge \bigwedge_{j \in S_1} \bar{A}_j \mid \bigwedge_{j \in S_2} \bar{A}_j \right]}{\Pr \left[ \bigwedge_{j \in S_1} \bar{A}_j \mid \bigwedge_{j \in S_2} \bar{A}_j \right]} \\
&\leq \frac{\Pr [A_i]}{e^{-1}} \leq p \cdot e \leq \frac{1}{d+1}
\end{aligned}$$

This concludes the proof of the lemma. ■

We will now apply the lemma to prove the theorem.

**Proof** We need to show that desired event occurs with positive probability. This probability can be bounded by

$$\begin{aligned}
\Pr \left[ \bigwedge_{i=1}^n \bar{A}_i \right] &= \Pr \left[ \bar{A}_1 \mid \bar{A}_2 \wedge \bar{A}_3 \wedge \dots \wedge \bar{A}_n \right] \\
&\quad \cdot \Pr \left[ \bar{A}_2 \mid \bar{A}_3 \wedge \bar{A}_4 \wedge \dots \wedge \bar{A}_n \right] \\
&\quad \vdots \\
&\quad \cdot \Pr \left[ \bar{A}_n \right] \\
&\geq \left( 1 - \frac{1}{d+1} \right)^n > 0
\end{aligned}$$

where the inequality follows from lemma 3. ■

## 2.1 Application: $k$ -SAT

We use the Lovász Local Lemma to prove that any  $k$ -CNF formula is satisfiable if a certain condition holds. First we define the  $k$ -SAT problem:

**Input** Boolean formula  $\varphi = \bigwedge_{i=1}^n C_i$  in  $k$ -CNF.

with  $m$  boolean variables  $x_1, \dots, x_m$

and  $n$  clauses  $C_1, \dots, C_n$  being conjunctions of  $k$  different literals

**Output** Decide whether there is a satisfying interpretation of  $\varphi$

We formalize the condition to guarantee the satisfiability of an  $k$ -CNF formula.

**Lemma 4** Any instance  $\varphi$  of  $k$ -SAT in which no variable appears in more than  $\frac{2^{k-2}}{k}$  clauses is satisfiable.

**Proof** We take an assignment  $X : \{x_1, \dots, x_m\} \rightarrow \{true, false\}$  uniformly at random. Let  $A_i$  be the "bad" event that clause  $C_i$  is unsatisfied by the random assignment  $X$ .

Let us determine the probability  $\Pr[A_i]$ . Since each clause consists of  $k$  different literals and each literal evaluates to false with probability  $\frac{1}{2}$  we get:

$$\Pr[A_i] = \frac{1}{2^k} =: p$$

Now let us bound the maximum dependency of  $A_i$ . Let  $j \neq i$ . It is obvious that  $A_i$  is dependent on  $A_j$  if and only if  $C_i$  and  $C_j$  share a variable. Each variable may occur in at most  $\frac{2^{k-2}}{k}$  clauses. Since  $C_i$  consists of  $k$  literals, it can be dependent on at most  $k \frac{2^{k-2}}{k} = 2^{k-2}$  other clauses.

Thus we can apply the Lovász Local Lemma with  $p = \frac{1}{2^k}$  and  $d = 2^{k-2}$  and get  $p \cdot (d+1) \cdot e \leq p \cdot d \cdot 4 = 1$  for sufficiently large  $d$ . Applying the Lovász Local Lemma we get that the probability of getting a satisfying assignment  $\Pr[\bigwedge_{i=1}^n A_i] > 0$ . Using the probabilistic method, we can conclude that there is in fact a satisfying assignment for  $\varphi$ . ■

## 2.2 Exercise 1: Application: 2-COLORING in a hyper graph

We use Lovász Local Lemma to prove that certain hyper graphs always have a 2-Coloring. A hyper graph is a tuple  $H = (V, E)$  with vertices  $V$  and hyper edges  $E$ . That is, each edge is a subset of the set of vertices  $V$ . A 2-Coloring is an assignment  $c : V \rightarrow \{red, blue\}$  of colors to vertices such that there is no monochromatic hyper edge. A monochromatic edge is an edge connecting only nodes of the same color.

**Exercise 1** Let  $H = (V, E)$  be a hyper graph in which every edge has at least  $k$  elements and intersects at most  $d$  other edges. For which  $k, d$  has  $H$  a 2-Coloring?

**Solution** We take a coloring uniformly at random. Let  $A_e$  be the bad event that edge  $e$  became monochromatic. This is the case when all vertices of  $e$  are either blue or all red. Now consider the probability of  $A_e$ :

$$\Pr[A_e] = \left(\frac{1}{2}\right)^k + \left(\frac{1}{2}\right)^k = 2^{1-k} =: p$$

We need to choose  $d$  such that the condition of the Lovász Local Lemma is satisfied. That is,  $p \cdot (d+1) \cdot e \leq 1$ . This can be achieved by choosing  $d \leq \frac{2^{1-k}}{e} - 1$ .

### 2.3 Exercise 2: Subtrees

The proof of Lovász Local Lemma given beforehand was non-constructive. Later on we will give a constructive proof. To do so, the following statement will be useful.


**Exercise 2** Consider a graph  $G$  of degree at most  $d + 2$ . Give an upper bound on the number of subtrees of  $G$  consisting of  $s$  nodes.

**Solution** For a fixed root we order the vertices in an arbitrarily (but fixed) order. Note, that this also orders the neighbors of a vertex uniquely. We will encode each tree with a bitstring of length  $(d + 1) \cdot s$  containing  $s - 1$  ones. We count the number of trees rooted at a vertex  $v$ .

Assume we had such an encoding. Then, the number of trees can be at most the number of bitstrings of this kind. That is:

$$\binom{(d+1) \cdot s}{s-1} \leq \left( \frac{(d+1) \cdot s \cdot e}{s-1} \right)^s \leq ((d+2) \cdot e)^s$$

It remains to be shown how a tree  $T$  can be encoded in such a fashion. The general idea is to do a pre-order traversal of  $T$  starting at the root  $v$ . When we discover a new vertex  $u$ , we output a string of length  $(d + 1)$ . This string indicates whether for every neighbor  $w$  of  $u$ ,  $w$  is contained in the tree. In particular, the string has a "1" at the  $i$ th position if and only if, the  $i$ th neighbor is a child of  $u$  in  $T$ . This is visualized in figure 2:



**Figure 2:** Tree of size 4 with coding 1100 0000 1000 0000

Because we have  $n$  possible roots, the total number of possible trees can be bounded by  $n \cdot ((d+2) \cdot e)^s$ .

## 2.4 A Constructive Proof of Lovász Local Lemma

The previous proof in this lecture was non-constructive. It did not show how to obtain an algorithm that computes the desired object. In this subsection we present the constructive version of this proof in [2] and [3] by giving a randomized algorithm for  $k$ -SAT. For the general case a similar algorithm can be constructed. We use the notation as before.

---

### Algorithm 1 Randomized $k$ -SAT


---

- 1: Pick random truth assignment  $X : \{x_1, \dots, x_m\} \rightarrow \{true, false\}$
  - 2: **while** there is an unsatisfied clause  $C$  **do**
  - 3:     Reflip all variables in  $C$
  - 4: **end while**
  - 5: Return the satisfying assignment
- 

Obviously this algorithm returns a satisfying assignment on termination. It remains to be shown that it in fact terminates in acceptable runtime.

**Theorem 5** *Let  $p \cdot (d + 1) \cdot e \leq 1 - \varepsilon$  for a constant  $\varepsilon > 0$ . Then the expected amount of reflips in the algorithm will be in  $\Theta(n)$ .*

**Proof** Let  $C_1, C_2, \dots$  be the (possibly infinite) sequence of clauses that are reflipped by the algorithm in that order. With each  $C_t$  we associate the tree  $T_t$  with a root node labeled as  $C_t$ . We add clause  $C_1, \dots, C_{t-1}$  in reversed order as follows: For  $i = t-1, \dots, 1$  we ignore  $C_i$  if it does not share any variable with any clause in  $T_t$ . Otherwise, we add  $C_i$  as a child of the deepest clause sharing at least one variable with  $C_t$ .



**Figure 3:** Example with reflipped clauses  $(x_1 \vee \bar{x}_3 \vee \bar{x}_4)$ ,  $(\bar{x}_1 \vee \bar{x}_2 \vee x_6)$ ,  $(x_3 \vee x_4 \vee \bar{x}_5)$  and  $(x_1 \vee \bar{x}_4 \vee x_5)$

We use the constructed tree  $T_t$  to prove the theorem by bounding the expected number of reflips. To do so, we have to measure the probability of a certain tree appearing.

**Claim** Consider any tree  $T_t$  of  $s$  nodes. The probability of this tree appearing is at most  $2^{-sk}$ .

**Proof** Consider a leaf of maximal depth in  $T_t$ . The corresponding clause does not share any variables with a clause which has been reflipped earlier. Thus, this clause was unsatisfied by the original assignment. This happens with probability  $2^{-k}$ . Since every variable in this clause is reflipped, the assignment of variables of higher clauses is independent. Hence, each node independently appears with probability  $2^{-k}$ . Because the tree consists of  $s$  nodes, its probability of appearing is  $2^{-sk}$ . ■

We use the claim to calculate the estimated amount of reflips for finding a satisfying assignment.

$$\begin{aligned}
\mathbb{E}[\#reflips] &= \mathbb{E}[\# \text{ trees appearing}] \\
&= \sum_{s=1}^{\infty} \mathbb{E}[\# \text{ trees of size } s] \\
&\leq n \cdot \sum_{s=1}^{\infty} \overbrace{((d+1) \cdot e)^s}^{\text{bound on the amount of trees}} \cdot \underbrace{2^{-sk}}_{\text{probability of a tree appearing}} \\
&= n \cdot \sum_{s=1}^{\infty} ((d+1) \cdot e \cdot 2^{-k})^s \\
&= n \cdot \sum_{s=1}^{\infty} ((d+1) \cdot e \cdot p)^s \\
&\leq n \cdot \sum_{s=1}^{\infty} (1 - \varepsilon)^s \\
&\leq \frac{n}{1 - (1 - \varepsilon)} = \frac{n}{\varepsilon} = \Theta(n)
\end{aligned}$$

This finishes the proof of the theorem. ■

## References

- [1] Paul Erdos and László Lovász. Problems and results on 3-chromatic hypergraphs and some related questions. *Infinite and finite sets*, 10:609–627, 1975.
- [2] Robin A Moser. A constructive proof of the lovász local lemma. In *Proceedings of the 41st annual ACM symposium on Theory of computing*, pages 343–350. ACM, 2009.
- [3] Robin A Moser and Gábor Tardos. A constructive proof of the general lovász local lemma. *Journal of the ACM (JACM)*, 57(2):11, 2010.