

Homework III, Topics in Theoretical Computer Science 2014

Due on Thursday April 17 at 14:00 (send an email to ola.svensson@epfl.ch or leave it in my letter box outside my office INJ 112). Solutions to many homework problems, including problems on this set, are available on the Internet, either in exactly the same formulation or with some minor perturbation. It is *not acceptable* to copy such solutions. It is hard to make strict rules on what information from the Internet you may use and hence whenever in doubt contact Ola Svensson. You are, however, allowed to discuss problems in groups with up to three students, but solutions should be handed in individually and please write with whom you have collaborated.

- 1 (70 pts) Johan Håstad proved that there is a particularly simple PCP verifier for SAT. For every $\epsilon > 0$ it uses the $O(\log n)$ random bits to compute three positions in the proof, say i, j , and k , and a bit b , and accepts iff

$$\pi(i) + \pi(j) + \pi(k) = b \pmod{2}.$$

Here $\pi(i)$ is the i 'th bit in the proof π . The verifier has completeness $1 - \epsilon$ and soundness $1/2 + \epsilon$. In other words,

- If φ is a satisfiable SAT instance then there is a proof π that makes the verifier accept with probability at least $1 - \epsilon$.
- If φ is a not satisfiable SAT instance then for any proof the verifier accepts with probability at most $1/2 + \epsilon$.

Your task in this problem is to use the above described verifier to prove the following statements:

- 1a (15 pts) For any $\epsilon > 0$, it is NP-hard to approximate MAX-3LIN within a factor of $1/2 + \epsilon$. MAX-3LIN is the problem where the input is a system of linear equations (modulo 2); each equation contains at most 3 variables; and we wish to find an assignment to the variables that satisfies the maximum number of equations.
- 1b (20 pts) For any $\epsilon > 0$, it is NP-hard to approximate MAX-3SAT within a factor $7/8 + \epsilon$. (*Hint: Do a reduction directly from MAX-3LIN.*)

- 1c** (35 pts) It is NP-hard to approximate the Vertex Cover problem within a factor of c , for some c that you wish to maximize.

For full score, show that c can be chosen to be $7/6 - \epsilon$ for any $\epsilon > 0$.¹

Recall that in the Vertex Cover problem we are given an undirected graph $G = (V, E)$ and we wish to find a subset $V' \subseteq V$ of minimum cardinality such that each edge is covered. An edge $\{u, v\} \in E$ is covered if $u \in V'$ or $v \in V'$.

- 2** (30 pts, Simplified Problem 11.8 in the book of Arora & Barak) Show that if $\text{SAT} \in \mathbf{PCP}(\log \log n, 1)$ then $\mathbf{P} = \mathbf{NP}$.

(Hint: What is the length of a proof for a $\mathbf{PCP}(r(n), 1)$ verifier? Recall that we assume that the proof only contains bits that are read with non-zero probability.)

BONUS PROBLEM

- 3** (bonus 30 pts, Problem 11.8 in the book of Arora & Barak) Show that if $\text{SAT} \in \mathbf{PCP}(r(n), 1)$ for some $r(n) = o(\log n)$ then $\mathbf{P} = \mathbf{NP}$. This shows that the PCP theorem is probably optimal up to constants.

¹This may be a little bit difficult but you will get partial credit for any constant $c > 1$ and the larger c the more credit.